

New Horizons LEISA KEM2 Cruise

Calibrated Data Overview

Abstract

This collection contains a set of calibrated images in FITS format taken by the Ralph Linear Etalon Imaging Spectral Array (LEISA) instrument during New Horizons' second Kuiper belt extended mission (KEM2).

Data Set Overview

This data set contains Calibrated data taken by the New Horizons Linear Etalon Imaging Spectral Array (LEISA) instrument during the KEM2 ENCOUNTER mission phase.

LEISA is an infrared imaging spectrometer. The two-dimensional (2-D) detector is a 256x256-pixel array. Spectral separation is done with a wedged optical etalon filter, which comprises high (2.1-2.25 micron) and low (1.25-2.5 micron) spectral resolution segments. The optics and filter arrangement configure one detector dimension as spatial at constant wavelength and the other dimension as both spectral and spatial. Multiple frames (images) may be taken in sequence, while scanning the Field Of View (FOV) in the spatial+spectral dimension, such that each pixel-sized piece of the observed target will be observed at all wavelengths across multiple frames. The resulting data products in this data set are three-dimensional image cubes, which dimensions are spatial, spatial+spectral, and spatial+time and which shall be convolved into spatial, spatial, and spectral image cubes at a later date.

Version History

Each subsection below details the major changes between the prior versions of this data set, listing the newest versions before older versions.

PDS4 v1.0

This version includes data acquired by the spacecraft between 10/01/2022 and 01/01/2026. It only includes data downlinked before 12/31/2025. Future datasets may include more data acquired by the spacecraft after 10/01/2022 but downlinked after 01/01/2026.

This version provides the results of new functional tests for LEISA.

This dataset corresponds to New Horizons NAIF SPICE distribution v0010.

General statement about data set versions after V1.0

Data files in versions after PDS4 v1.0 will not be reprocessed if the only updates are normal SPICE improvements.

Processing

The data in this data set were created by a software data processing pipeline on the Science Operations Center (SOC) at the Southwest Research Institute (SwRI), Department of Space Operations. This SOC pipeline assembled data as FITS files from raw telemetry packets sent down by the spacecraft and populated the data labels with housekeeping and engineering values, and computed geometry parameters using SPICE kernels. The pipeline did not resample the data.

Calibration

Detailed information about calibration of LEISA data is available in the SOC Instrument Interface Control Document (ICD) in the DOCUMENT section of this data set. The LEISA calibration will only be briefly summarized here; refer to the ICD for details about what is summarized here.

The calibration of LEISA images comprises the following steps:

- 1) Remove electronics-induced and flat-field signal
- 2) Apply calibration offset and gain
- 3) Adjust for integration time, filter width, and pixel solid angle
- 4) Correct for gain

In addition, the calibration procedure calculates various quantities such as error (see note below) and a data quality flag for each pixel and includes those results in the calibrated data product as additional PDS OBJECTs (FITS extensions) appended to the main OBJECT with the data image. The quality flag PDS OBJECT is an image of values of the same size as the main IMAGE product, with each quality flag pixel mapped to the corresponding pixel in the main product. A quality flag value of zero indicates a valid pixel; a non-zero value indicates an invalid pixel.

Note that for windowed products, all pixels in an image are not returned in the downlink telemetry. In the raw data, the pipeline sets such pixels to zero DN (Data Number); the calibration processes those zero-DN pixels as if they were real raw values may result in a confusing result with the majority of the displayed image appearing as an inverse of the calibration (calibration of zero values); therefore the windowed status of the image, as recorded in the value for the NOTE keyword of the PDS label should be considered when looking at these data.

Calibration and data characteristics

- i) Error estimates are all zero. As of late 2016, the code to calculate the values for the Error estimates extension has not been deployed to the SOC, and placeholder code sets all pixel error estimates to zero. When the updated code is deployed to the SOC, some reprocessed observations may be delivered to PDS (e.g. Jupiter encounter data); note that observations from the Launch and Pluto Cruise mission phases have little if any science utility, so there is no incentive to reprocess those data just to provide error estimates.

- ii) Fixed-pattern noise. There is usually fixed-pattern noise (FPN) in the data that cannot be removed by the flat-field correction. Users need to generate an average of a few run-up frames containing background sky and the FPN, and subtract that average from every frame with a target source present. The number of frames to use will vary by observation, so this operation is not part of the automated pipeline.
- iii) Scattered light. Although some observations have been taken to characterize scattered light, no photometric modeling of the stray light has been made or applied to the data. Those observations have only been used to optimize the planned imaging at Pluto encounter to minimize the effect of stray light.

Ongoing in-flight calibration observations will be analyzed to assess the long term stability of the calibration.

Data

The observations in this data set are stored in data files using standard Flexible Image Transport System (FITS) format. Each FITS file has a corresponding detached PDS label file, named according to a common convention. The FITS files may have image and/or table extensions. See the PDS label plus the document collection for a description of these extensions and their contents.

This Data section comprises the following sub-topics:

- Filename/Product IDs
- Instrument description
- Other sources of information useful in interpreting these Data
- Visit Description, Visit Number, and Target in the Data Labels

Filename/Product IDs

The filenames and Local product Identifiers (LID) of observations adhere to a common convention, e.g.:

ApIDs	Data product description/Prefix(es)
0x53c	LEISA Lossless (CDH 1)/LRW,LSB
0x54b	LEISA Lossless (CDH 2)/LRW,LSB
0x53d	LEISA Packetized (CDH 1)/LRW,LSB
0x54c	LEISA Packetized (CDH 2)/LRW,LSB
0x53e	LEISA Lossy (CDH 1)/LRW,LSB
0x54d	LEISA Lossy (CDH 2)/LRW,LSB

There are other ApIDs that contain housekeeping values and other values. See the SOC Instrument ICD for more details: [urn:nasa:pds:nh_documents:mission:soc_inst_icd](#)

Please note that not all ApIDs may be found in this data set.

Instrument description

Refer to the following files for a description of this instrument.:

- New Horizon LEISA instrument overview:
[urn:nasa:pds:nh_documents:ralph:leisa_inst_overview](#)
- Ralph Space Science Review (SSR) paper:
[urn:nasa:pds:nh_documents:ralph:ralph_ssr](#)
- SOC Instrument ICD: [urn:nasa:pds:nh_documents:mission:soc_inst_icd](#)
- Ralph SPICE Instrument Kernel:
[urn:nasa:pds:nh_documents:ralph:nh_ralph_v100_ti](#)

Other sources of information useful in interpreting these Data

Refer to the following files for more information about these data:

- NH Mission Trajectory Table:
[urn:nasa:pds:nh_documents:mission:nh_mission_trajectory](#)
- Field of View Illustration: [urn:nasa:pds:nh_documents:mission:nh_fov](#)
- Ralph SPICE Instrument Kernel:
[urn:nasa:pds:nh_documents:ralph:nh_ralph_v100_ti](#)

Visit Description, Visit Number, and Target in the Data Labels

The observation sequences were defined in Science Activity Planning (SAP) documents and grouped by Visit Description and Visit Number. The SAPs are spreadsheets with one Visit Description & Number per row. A nominal target is also included on each row and included in the data labels but does not always match with the target name field's value in the data labels. In some cases, the target was designated as `right_ascension_angle`, `declination_angle` pointing values in the form "`right_ascension_angle`, `declination_angle` =123.45,-12.34" indicating Right Ascension and Declination, in degrees, of the target from the spacecraft in the Earth Equatorial J2000 inertial reference frame. This indicates that either the target was a star, or the target's ephemeris was not loaded into the spacecraft's attitude and control system which in turn meant the spacecraft could not be pointed at the target by a body identifier and an inertial pointing value had to be specified as Right Ascension and Declination values. PDS-SBN practices do not allow putting a value like `right_ascension_angle`, `declination_angle` =... in the PDS target name keyword's value. In those cases, the PDS target purpose value is set calibration. Target

name may be None for a few observations in this data set; typically, that means the observation is a functional test so None is an appropriate entry for those targets, but the PDS user should also check the `nh:observation_description` and `nh:sequence_id` keywords in the PDS label, plus the provided sequence list (`urn:nasa:pds:nh_documents:ralph:seq_leisa_kem2`) to assess the possibility that there was an intended target. These two keywords are especially useful for star targets as often stars are used as part of instrument calibrations and are included as part of the sequencing description which is captured in these keywords.

Ancillary Data

The geometry items included in the data labels were computed using the SPICE kernels archived in the New Horizons SPICE data set, NH-J/P/SS-SPICE-6-V1.0.

Every observation provided in this data set was taken as a part of a particular sequence. A list of these sequences has been provided within the NH Ralph document collection (PDS4 LID `urn:nasa:pds:nh_documents:ralph`) within the PDS, one file for each mission phase. The sequence identifier and description are included in the PDS label for every observation.

N.B. While every observation has an associated sequence, every sequence may not have associated observations. Some sequences may have failed to execute due to spacecraft events (e.g., safing). No attempt has been made during the preparation of this data set to identify such empty sequences.

Time

There are several time systems, or units, in use in this dataset: New Horizons spacecraft MET (Mission Event Time or Mission Elapsed Time), UTC (Coordinated Universal Time), and TDB (Barycentric Dynamical Time).

This section will give a summary description of the relationship between these time systems. For a complete explanation of these time systems the reader is referred to the documentation distributed with the Navigation and Ancillary Information Facility (NAIF) SPICE toolkit from the PDS NAIF node, (see <http://naif.jpl.nasa.gov/>).

The most common time unit associated with the data is the spacecraft MET. MET is a 32-bit counter on the New Horizons spacecraft that runs at a rate of about one increment per second starting from a value of zero at “19.January, 2006 18:08:02 UTC” or “JD2453755.256337 TDB.”

The leapsecond adjustment ($\Delta ET = ET - UTC$) was 65.184s at NH launch, and the first four additional leapseconds occurred at the ends of 12/2009, 06/2012, 06/2015, and 12/2016. Refer to the NH SPICE data set, NH-J/P/SS-SPICE-6-V1.0, and the SPICE toolkit documentation, for more details about leapseconds.

The data labels for any given product in this dataset usually contain at least one pair of common UTC and MET representations of the time at the middle of the observation. Other portions of the products, for example tables of data taken over periods of up to a day or more, will only have the MET time associated with a given row of the table.

For the data user's use in interpreting these times, a reasonable approximation (+/- 1s) of the conversion between Julian Day (TDB) and MET is as follows:

$$\text{JD TDB} = 2453755.256337 + (\text{MET} / 86399.9998693)$$

For more accurate calculations the reader is referred to the NAIF/SPICE documentation as mentioned above.

Reference Frame

Geometric Parameter Reference Frame

Earth Mean Equator and Vernal Equinox of J2000 (EMEJ2000) is the inertial reference frame used to specify observational geometry items provided in the data labels. Geometric parameters are based on best available SPICE data at time of data creation.

Epoch of Geometric Parameters

All geometric parameters provided in the data labels were computed at the epoch midway between the start_date_time and stop_date_time label fields.

Software

The observations in this data set are in standard FITS format with PDS labels and can be viewed by a number of PDS-provided and commercial programs. For this reason, no special software is provided with this data set.

Confidence Level Overview

During the processing of the data in preparation for delivery with this volume, the packet data associated with each observation were used only if they passed a rigorous verification process including standard checksums.

In addition, raw (CODMAC Level 2) observation data for which adequate contemporary housekeeping and other ancillary data are not available may not be reduced to calibrated (CODMAC Level 3) data. This issue is raised here to explain why some data products in the raw data set may not have corresponding data products in the calibrated data set.

Data coverage and quality

Every observation provided in this data set was taken as a part of a particular sequence. For this data set, these KEM2 sequences can be found in the ralph document collection under PDS4 LID `urn:nasa:pds:nh_documents:ralph:seq_leisa_kem2`. Please note that some sequences provided may have zero corresponding observations.

Refer to the Confidence Level Overview section above for a summary of steps taken to assure data quality.

Caveat about target name in PDS labels and observational

The downlink team on New Horizons has created an automated system to take various uplink products, decode things like Chebyshev polynomials in command sequences representing celestial body ephemerides for use on the spacecraft to control pointing, and infer from those data what the most likely intended target was at any time during the mission. This works well during flyby encounters and less so during cruise phases and hibernation.

The user of these PDS data needs to be cautious when using the target name and other target-related parameters stored in this data set. This is less an issue for the plasma and particle instruments, more so for pointed instruments. To this end, the heliocentric ephemeris of the spacecraft, the spacecraft-relative ephemeris of the inferred target, and the inertial attitude of the instrument reference frame are provided with all data, in the J2000 inertial reference frame, so the user can check where that target is in the Field Of View (FOV) of the instrument.

Finally, note that, within the FITS headers of the data products, the sequence tables, and other NH Project-internal documents used in this data set, informal names are often used for targets instead of the canonical names used within the PDS labels. For example, during the Pluto mission phase, instead of the target name '15810 ARAWN (1994 JR1)' there might be found any of the following: 1994JR1; 1994 JR1; JR1. However, within the context of this data set, these project abbreviations are not ambiguous (e.g. there is only one NH target with 'JR1' in its name), so there has been, and will be, no attempt to expand such abbreviations where they occur outside formal PDS keyword values.

Contact Information

For any questions regarding the data format of the archive, contact the New Horizons LEISA Principal Investigator: Alan Stern, Southwest Research Institute

S. Alan Stern
Southwest Research Institute (SwRI)
Department of Space Studies
1301 Walnut Street
Boulder, CO 80302
USA

Further Reading

Steffl, A.J., J. Peterson, B. Carcich, L. Nguyen, and S.A. Stern, NEW HORIZONS SPICE KERNELS, V1.0, NH-J/P/SS-SPICE-6-V1.0, NASA Planetary Data System, 2007.

<https://doi.org/10.17189/1520109>